



COMPARISON OF COLOUR VISION IN DIFFERENT TYPES OF DEVIATION

¹*Partha Haradhan Chowdhury and ²Brinda Haren Shah

¹Department of Optometry, Shree Satchandi Jankalyan Samiti Netra Prasikshan Sansthan, Pauri, Affiliated to Uttarakhand State Medical Faculty, Dehradun, India

²B.Optom, Department of Optometry, Gujarat University, Gujarat, India

ARTICLE INFO

Article History:

Received 28th March, 2018
Received in revised form
19th April, 2018
Accepted 24th May, 2018
Published online 30th June, 2018

Key Words:

Colour vision,
Esotropia,
Exotropia.

ABSTRACT

Purpose: The aim of the present study is to compare the colour vision in different types of deviation.

Methods: A pilot, cross sectional, observational study was performed at tertiary eye care centers. Subjects with Ocular deviation between 10 to 40 prism diopters, Corrected distance Colour vision should be greater than 6/18 and Age should be between 10 to 40 years of age were included in the study. Colour vision was assessed with Fransworth D 15 colour vision test.

Results: 30 subjects were included in the study. Out of that, 16 subjects were in the age group of 11-20 years, 12 subjects were in the age group of 21-30 years and 2 subjects were in the age group of 31-40 years. 60% subjects were Female and 40% subjects were Male. The mean colour vision was considered in each type of deviation. It shows that colour vision will be deteriorated more in Esotropia as compared to Exotropia

Conclusions: Deterioration of colour vision is observed more in Esotropia as compared to Exotropia.

Copyright © 2018, Partha Haradhan Chowdhury and Brinda Haren Shah. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Partha Haradhan Chowdhury and Brinda Haren Shah, 2018. "Comparison of colour vision in different types of deviation", *International Journal of Development Research*, 8, (05), xxxxxxxxxx.

INTRODUCTION

According to increasing ocular deviation CV is being deteriorated due to the anatomical consideration is the main factor. Increasing ocular deviation proportionately image is shifted from the foveal region and thus deterioration is present. In case of Esodeviation images are shifted to nasal fovea and due to very less time of intermittent stages are present in Eso deviation there may be lots of chances to become Amblyopia compare to Exo deviation. If Eso deviation is present during the visual development stages on that time it become very crucial to treat it specially in case of Eso deviation due to most of the time deviated eye is become fixed but in case of Exo deviation most of the time it is associated with intermittent deviation. That's why there may be lots of chances of deterioration of CS is more in Eso deviation compare to Exo deviation.

*Corresponding author: Partha Haradhan Chowdhury

Department of Optometry, Shree Satchandi Jankalyan Samiti Netra Prasikshan Sansthan, Pauri, Affiliated to Uttarakhand State Medical Faculty, Dehradun, India


But in case CV there may not be much more difference in deterioration between Eso and Exo deviation.

MATERIALS AND METHODS


A pilot, cross sectional, observational study was performed at tertiary eye care centers. Subjects with Ocular deviation between 10 to 40 prism diopters, Corrected distance Colour vision should be greater than 6/18 and Age should be between 10 to 40 years of age were included in the study. Individuals with any other systemic disease (specially which can affect study), Individuals with any other Ocular Pathology, with any active ocular infection, any ocular anomalies like Corneal Scar etc, ocular deviation if less than 10 degree and Significant amount of amblyopic patient were excluded from the study. Full refractive correction along with detailed fundus evaluation was performed in each and every patient. Colour vision was assessed with Fransworth D 15 colour vision test in different types of ocular deviation. Data was analyzed using SPSS software version 20.

RESULTS


30 subjects were included in the study. Graph 1 shows distribution of subjects in various age groups. 16 subjects were in the age group of 11-20 years, 12 subjects were in the age group of 21-30 years and 2 subjects were in the age group of 31-40 years. Graph 2 shows gender wise distribution of the subjects. 60% subjects were Female and 40% subjects were Male. Graph 3 shows comparison of Colour vision for different types of ocular deviation. The mean colour vision was considered in each type of deviation. It shows that colour vision will be deteriorated more in esotropia as compared to exotropia.



Graph 1. Shows age wise distribution of the subject



Graph 2. Shows gender wise distribution of the subjects



Graph 3. Shows comparison of Colour vision for different types of deviation

DISCUSSION

According to the present study, it has been showed that in cases of ocular deviation, images of an object fall on the parafoveal region and deteriorates the colour vision. Esodeviation is more harmful as compared to Exodeviation, because intermittent timings are very less in Esodeviation as compared to Exodeviation. In case of Esodeviation images of an object is fall on the nasal parafoveal region and according to the deformity of the anatomical arrangement of the photoreceptor cells in the macula that's why in the parafoveal region number of cone cells is less compare to foveal region. Just for this reason in case of Esodeviation images is shifted towards parafoveal region and due to less number of cone cells and according to the statistics it has been proved that with increasing Esodeviation color vision is been deteriorated. According to the study it has been proved that in case of Esodeviation the deterioration of color vision is more compare to Exodeviation.

Conclusion

Deterioration of colour vision is observed more in Esotropia as compared to Exotropia.

Consent: Oral/ written consent was obtained from patient as well as from tertiary eye care centres.

Ethical Approval: It is not applicable.

Competing Interests: Authors have declared that no competing interests exist.

REFERENCES

- A.G. KOCAK-ALTINTAS *et al* 2000. "Colour vision and Colour Vision deficiency in Amblyopia" *European Journal of Ophthalmology*, vol. 10, no.1, pg 77-81
- Alan W. Freeman *et al*. 1996. "Components of Colour vision Loss in Strabismus" published in the journal of vision research, Vol. 36, No. 5, Pg. 765-774
- Anika K. Tandon *et al*. 2014. "Binocular Inhibition in Strabismic Patients is Associated with Diminished Quality of Life", *Journal of American Association for Pediatric Ophthalmology and Strabismus*, October volume 18, issue 5, pg – 423-426
- Handbook of pediatric Strabismus and amblyopia, Kenneth W. Wright, Peter H. Spiegel, Lisa Thompson, First ed, 2006
- Hui Zhu, *et al*. 2015. "Association between Childhood Strabismus and Refractive Error in Chinese Preschool Children" *Journal of Plos One*.
- Ye, *et al*, 2014. "Strabismus genetics across a spectrum of eye misalignment disorders", *Journal of clinical genetics*, 2014, vole 86, pg 103-111
- Zhale Rajavi, *et al*, 2015. "Prevalence of Colour Vision Deficiency and its Correlation with Amblyopia and Refractive Errors among Primary School Children", *Journal of Ophthalmic and Vision Research*, vol. 10, issue 2, pg 130-138.